

The πN Sigma term – a lattice investigation^{1 2}

R. Altmeyer^a, M. Göckeler^b, R. Horsley^{b,a}, E. Laermann^c and G. Schierholz^{a,b}^aDESY, Notkestraße 85, D-22603 Hamburg, Germany^bHLRZ, c/o Forschungszentrum Jülich, D-52425 Jülich, Germany^cFakultät für Physik, Universität Bielefeld, D-33501 Bielefeld, Germany

Abstract

A lattice calculation of the πN sigma term is described using dynamical staggered fermions. Preliminary results give a sea term comparable in magnitude to the valence term.

Introduction and Theoretical Discussion

Hadrons appear to be far more complicated than the (rather successful) constituent quark model would suggest. For example a recent result from the EMC experiment, [1], suggests that constituent quarks are responsible for very little if any of the nucleon spin. Another, older result, is from the πN sigma term, see e.g. [2], which seems to give a large strange component to the nucleon mass. To explain theoretically these results and other experiments involves the computation of certain matrix elements – a non-perturbative calculation. This is thus an area where lattice calculations may be of some help.

The πN sigma term, $\sigma_{\pi N}$, is defined³ as that part of the mass of the nucleon (for definiteness the proton) coming from the vacuum connected expectation value of the up (u) and down (d) quark mass terms in the QCD Hamiltonian,

$$\sigma_{\pi N} = m \langle N | \bar{u}u + \bar{d}d | N \rangle, \quad (1)$$

where we have taken these quarks to have equal current mass ($= m$). Other contributions to the nucleon mass come from the chromo-electric and chromo-magnetic gluon pieces and the sea terms due to the s quarks. Experimentally this matrix element has been measured from low energy π - N scattering. A delicate extrapolation to the chiral limit [3, 4] gives a result for the isospin even amplitude of Σ/f_π^2 with $\Sigma = \sigma_{\pi N}$, from which the πN sigma term may be found. The precise value obtained this way has been under discussion for many years. However within the limits of our lattice calculation, this will not concern us here and for orientation we shall just quote a range of results from later analyses of $\sigma_{\pi N} \approx 56\text{MeV}$, [5], down to 45MeV , [6].

To estimate valence and sea contributions to $\sigma_{\pi N}$, classical current algebra analyses assume octet dominance and make first order perturbation theory about the $SU_F(3)$ flavour symmetric Hamiltonian. We have

$$\hat{\mathcal{H}}_m = m(\bar{u}u + \bar{d}d) + m_s \bar{s}s \equiv m_{av} \bar{\psi} I \psi - \frac{m_s - m}{\sqrt{3}} \bar{\psi} \lambda^8 \psi. \quad (2)$$

The first term in the second equation is the flavour symmetric Hamiltonian with mass $m_{av} = (2m + m_s)/\sqrt{3}$, while the second term is the flavour breaking piece. (ψ is the column vector (u, d, s) .) We thus have from eq. (1)

$$\begin{aligned} \sigma_{\pi N} &\approx m \langle N | \bar{u}u + \bar{d}d - 2\bar{s}s | N \rangle_{symm} + 2m \langle N | \bar{s}s | N \rangle_{symm} \\ &\stackrel{def}{=} \sigma_{\pi N}^{val} + \sigma_{\pi N}^{sea}, \end{aligned} \quad (3)$$

where we have first assumed that the nucleon wavefunction does not change much around the symmetric point. We then subtract and add a strange component. At the symmetric point the u and d quarks each have equal valence and sea part, while the s quark matrix element only has a sea component. Thus in the first term the sea contribution cancels, justifying the definitions given in the last line of eq. (3). Also this means that in the following we shall interchangeably talk about ‘strange’ and ‘sea’ contributions to

¹The MT_c Collaboration.

²Talk presented by R. Horsley at the Workshop on Field-Theoretical Aspects of Particle Physics, Kyffhäuser, Germany, September 13th to 17th 1993. To be published in the NTZ Series.

³Actually, originally as the matrix element of the double commutator of the Hamiltonian with two axial charges. However this is equivalent to the definition given in eq. (1), see for example ref. [3].

the nucleon. At first order perturbation theory $\sigma_{\pi N}^{val}$ may easily be calculated⁴ to give $\sigma_{\pi N}^{val} \approx 25\text{MeV}$ and so $\sigma_{\pi N}^{sea} \approx 31 \sim 20\text{MeV}$. This in turn means that

$$m_s \langle N | \bar{s}s | N \rangle \approx 400 \sim 250\text{MeV}, \quad (4)$$

which would indicate a sizeable portion of the nucleon mass comes from the strange quark contribution. (Remember that the mass of the proton is about 938MeV.) This would not be expected from the constituent quark model, where the nucleon is made only of (dressed) u and d quarks.

Staggered Fermions

As we see from above, we need to calculate certain matrix elements. One of the most promising methods at present to calculate them comes from the lattice, where Euclidean space-time is discretised with a lattice spacing a . The problem is then turned into a statistical mechanical one of evaluation of correlation functions (as to be described below). This can be attacked using Monte Carlo simulations. The continuum limit is obtained when $a \rightarrow 0$, or $\beta \equiv 6/g^2(a) \rightarrow \infty$. This procedure is well known, see for example [8]. We have performed a large scale simulation on a $16^3 \times 24$ lattice of QCD using the Hybrid Monte Carlo method. The main emphasis was for $\beta = 5.35$, $m = 0.01$, although some other runs were also performed. (Our results and further details are given in [9].)

There are two commonly used formulations of fermions on the lattice – Wilson and staggered. Both have their advantages and disadvantages. We have used the staggered approach, which describes in the continuum 4 degenerate flavours. Here the fermion fields in the partition function, $\chi^a(x)$ are only a function of the lattice point and colour index. Quark spin and flavour combinations are given by summing appropriately over χ 's sitting on the vertices of hypercubes, thus

$$q_{\alpha f}(X) = \sum_A \Gamma_{\alpha f}^A \chi(X + A). \quad (5)$$

The hypercube is defined by $x = X + A$, A_μ being 0 or 1 and X_μ even, where x is the lattice site. Γ^A is given by $\gamma_1^{A_1} \dots \gamma_4^{A_4}$. Eq. (5) projects out the flavour (f) and spin (α) degrees of freedom. In the continuum limit we have for $m = 0$ a chiral symmetry $SU_L(4) \otimes SU_R(4)$. The diagonal symmetry, valid for all m , is the flavour group $SU_F(4)$. The advantage of the staggered lattice fermion formulation is that we have, for $m = 0$, a remnant $U(1) \otimes U(1)$ symmetry. This allows a clean study of chiral symmetry breaking as $m \rightarrow 0$, and also the emergence of a massless π , with $m_\pi^2 \propto m$. (For Wilson fermions, one must carefully tune the hopping parameter to obtain a massless π , chiral properties always being explicitly broken.) As we saw in the last section we have at the symmetric point an $SU_F(3)$ symmetry for the u, d, s quarks. (Corrections for the strange quarks are treated in first order perturbation theory, as part of the sea contribution.) As we shall be here interested in the questions of sigma/nucleon and valence/sea ratios, we expect that the different number of quarks present will make little difference.

Lattice calculations have to be performed when the lattice spacing a is small enough that finite lattice effects are negligible. Staggered fermions have a natural indicator to show this by when the flavour symmetry is restored. On the lattice meson ($M \sim \bar{\chi}\chi$) and baryon ($B \sim \chi\chi\chi$) operators which in the continuum describe degenerate states lie in several distinct lattice representations, each with a different mass. As $a \rightarrow 0$ these masses must tend to a common mass. Practically we say that we are close enough to the continuum when possible mass differences are negligible. For our configurations we seem to be entering such a region, [9].

Another technical point concerns the existence of a parity partner in the correlation functions. Masses (M_α) and amplitudes ($A_{\alpha\alpha}$) are measured from 2-point correlation functions. For the baryon we have

$$C(t) \stackrel{def}{=} \langle B(t) \bar{B}_W(0) \rangle$$

⁴Baryon octet mass splittings are given by

$$M_B - M_{symm} = -(m_s - m) \langle B | \bar{\psi} \lambda^8 \psi | B \rangle_{symm} / \sqrt{3},$$

where B is the $\frac{1}{2}^+$ nucleon multiplet. From the Wigner-Eckart theorem for $SU(3)$ we have

$$\langle B | \bar{\psi} \lambda^8 \psi | B \rangle_{symm} = F \text{tr}(B^\dagger [\lambda^8, B]) + D \text{tr}(B^\dagger \{\lambda^8, B\}).$$

From the known baryon masses F, D can then be estimated. (We use the numbers given in [7] as $(m_s - m)F \approx 190$, $(m_s - m)D \approx -61$.) Furthermore

$$\sigma_{\pi N}^{val} = m\sqrt{3} \langle N | \bar{\psi} \lambda^8 \psi | N \rangle_{symm} = m(3F - D).$$

Using $m_s/m \approx 26$ gives the quoted results.

$$\begin{aligned}
&\approx \langle 0 | \hat{B} \hat{S}_4^t \hat{\bar{B}}_W | 0 \rangle \\
&\approx A_{NN} \mu_N^t + A_{\Lambda\Lambda} \mu_\Lambda^t \quad \frac{1}{2}T \gg t \gg 0.
\end{aligned} \tag{6}$$

In the second line we have gone from the (numerically) calculable 2-point correlation function to the equivalent expression in the operator formalism. The transfer matrix is denoted by $\hat{T} = \hat{S}_4^2 \equiv \exp(-2\hat{H})$. A shift of one unit still lies in the hypercube, which then involves a flavour transformation. Thus we can write $\hat{S}_4 = \hat{\Xi}_4 \hat{T}^{\frac{1}{2}}$ where $\hat{\Xi}_4$ is a non-local flavour transformation. (No explicit representation of $\hat{\Xi}_4$ is at present known.) Eigenstates of $\hat{S}_4$ are defined by

$$\hat{S}_4 |\alpha\rangle = \mu_\alpha |\alpha\rangle \quad \mu_\alpha = \xi_\alpha e^{-M_\alpha}, \tag{7}$$

($\langle \alpha | \alpha \rangle = 1$) where $\xi_\alpha = \pm 1$ is the eigenvalue of $\hat{\Xi}_4$. Inserting complete sets of states in the second line of eq. (6) gives the third line where the amplitude $A_{\alpha\beta} = \langle 0 | \hat{B} | \alpha \rangle \langle \beta | \hat{\bar{B}}_W | 0 \rangle$, and $\mu_N = \exp(-M_N)$, $\mu_\Lambda = -\exp(-M_\Lambda)$. (We shall call Λ , a little artificially, the parity partner to the nucleon.) The time box length (T , here = 24) has been assumed to be so large that there are no finite time-size effects (although these can be easily taken into account). The parity partner arises because on the lattice inversion $I_s : x_i \rightarrow -x_i$ is not parity (P) as it effects the flavour degrees of freedom. (One can show that $\hat{P} = \hat{I}_s \hat{\Xi}_4$.) As a consequence hadron operators in general do not have a definite parity.

As an example of a correlation function we show in Fig. 1 a baryon correlation function. The typical




Figure 1: A typical baryon correlation function plotted against t for $\beta = 5.35$, $m = 0.01$. Boxes denote negative values. The fit interval is from $t = 2$ to 22 .

zigzagging of the results (and fit) is caused by the parity partner. To pick out the ground states we must take $t \rightarrow \infty$ and/or choose an operator with a good overlap with this state. Experience has shown that a non-local operator (the ‘wall’ and hence the subscript ‘ W ’) has rather good overlap properties with the ground state for very moderate t , so this has been used as the source; the sink is taken as a local baryon operator. The mass obtained this way agrees with that which comes from using a local baryon operator for both sink and source.

Measuring $\sigma_{\pi N}$

Practically there are several possibilities open to us for the evaluation of the matrix element. The easiest is simply to differentiate eq. (7). As⁵ $\frac{\partial \hat{S}_4}{\partial m} = -\bar{\chi} \chi \hat{S}_4$ we have

$$m \frac{\partial M_N}{\partial m} |_\beta = \langle N | m \bar{\chi} \chi | N \rangle = \sigma_{\pi N}, \tag{8}$$

an application of the Feynman-Hellmann theorem. Thus all we need to do is to measure M_N for different masses m (at constant β) and numerically estimate the gradient. In Fig. 2 we show such a procedure. (At present we have simply fitted a straight line through all points.) We find

$$\sigma_{\pi N} \approx 11.9(8)m|_{m=0.01} \approx 0.12(1), \tag{9}$$

⁵In the staggered formalism the operator expression for $m \bar{\chi} \chi$ has a more complicated appearance than that given here, [9, 10].




Figure 2: The Nucleon mass, M_N , plotted against m for $\beta = 5.35$.

which gives

$$\frac{\sigma_{\pi N}}{M_N} \approx \frac{11.9m}{11.9m + 0.64} \Big|_{m=0.01} \approx 0.15, \quad (10)$$

which is to be compared with the experimental result of $\sigma_{\pi N}/M_N \approx 0.06 \sim 0.05$. The numerical result is much larger, but presumably this simply indicates that we have used much too large a quark mass in our simulation (which at present we, and everybody else, cannot avoid doing due to computer limitations).

We now wish to estimate the valence and sea contributions. This is technically more complicated and involves the evaluation of a 3-point function, [11]

$$C(t; \tau) = \langle B(t) m \bar{\chi} \chi(\tau) \bar{B}_W(0) \rangle, \quad (11)$$

This may be diagrammatically sketched as the sum of two terms:




Diagram (a) is the connected (or valence) piece; the nucleon quark lines are connected with the mass insertion operator. Diagram (b) represents the disconnected (or sea) piece.

For the connected piece we have fixed t to be 8 or 9 and then evaluated $C(t; \tau)$ as a function of τ . The appropriate fit function is

$$C(t; \tau) \approx \sum_{\alpha, \beta=N, \Lambda} A_{\alpha\beta} \langle \alpha | m \bar{\chi} \chi | \beta \rangle \mu_{\alpha}^{t-\tau} \mu_{\beta}^{\tau} \quad \frac{1}{2}T \gg t \gg \tau \gg 0. \quad (12)$$

The $A_{\alpha\beta}$ and M_{α} are known from 2-point correlation functions. We see that when $\alpha = \beta = N$ we have the matrix element that we require: $\langle N | m \bar{\chi} \chi | N \rangle$. However there are other terms which complicate the fit: $\langle \Lambda | m \bar{\chi} \chi | \Lambda \rangle$ and the cross terms $\langle \Lambda | m \bar{\chi} \chi | N \rangle \equiv \mu_{\Lambda} \mu_N^{-1} \langle N | m \bar{\chi} \chi | \Lambda \rangle$. As can be seen from eq. (12) these cross terms are responsible for oscillations in the result. To disentangle the wanted result from $\langle \Lambda | m \bar{\chi} \chi | \Lambda \rangle$ we need to make a joint fit to the $t = 8$ and $t = 9$ results. At present we have not done this,




Figure 3: The ratio of the 3-point connected correlation function to the 2-point correlation function against τ for $\beta = 5.35$, $m = 0.01$. The points used in the fit are joined together. (The dotted lines just extend the fit to neighbouring points.)

but just checked that separate fits give consistent results. In Fig. 3 we show preliminary results for $t = 8$. Up until now we have only evaluated about a quarter of our available configurations, so the final results should have reduced error bars. We find

$$\sigma_{\pi N}^{val} \approx 0.08(2), \quad (13)$$

with about the same result for the Λ matrix element. The cross term is smaller, roughly 0.02.

Finally we have attempted to estimate the disconnected term using a stochastic estimator, [12]. This is costly in CPU time. One can improve the statistics by summing the 3-point correlation function over τ ; this is then equivalent to a differentiation of the 2-point function with respect to m . This gives

$$\sum_{\tau} C(t; \tau) \approx t \sum_{\alpha=N, \Lambda} A_{\alpha\alpha} \langle \alpha | m \bar{\chi} \chi | \alpha \rangle \mu_{\alpha}^t \quad \frac{1}{2}T \gg t \gg 0. \quad (14)$$

In Fig. 4 we show the disconnected part of the 3-point correlation function. As expected the quality of




Figure 4: The ratio of the 3-point disconnected correlation function to the 2-point correlation function against t . 100 sets of Gaussian random numbers were used for the stochastic estimator. To improve the signal we have averaged over the points t and $T - t$.

the data is poor – optimistically we see a slope. A simple linear fit gives

$$\sigma_{\pi N}^{sea} \approx 0.05(4). \quad (15)$$

Discussion

We see that (roughly at least) eq. (9) is consistent with eqs. (13,15) and that

$$\frac{\sigma_{\pi N}}{\sigma_{\pi N}^{val}} \approx 1.5, \quad (16)$$

which is to be compared with the experimental value of about $2.2 \sim 1.8$. Although we can draw no firm conclusions at present our result tentatively indicates that the valence part of the πN sigma term is slightly larger than the sea term.

Comparing our results with other work obtained using dynamical fermions, Gupta et. al, [7] who use 2 flavours of Wilson fermions, find $\sigma_{\pi N}/\sigma_{\pi N}^{val} \approx 2 \sim 3$, which indicates a somewhat larger sea component in the πN sigma term. On the other hand Patel, [13], using results from [14] for 2 staggered flavours finds for the ratio $1.5 \sim 2.0$, while Bernard et. al., [15], have ~ 2.0 . These, like our result, seems to be lower than for the Wilson fermion case.

We would also like to emphasise that although lattices can give a first principle calculation, at present one is not able to do this. Technically the fit formulæ that we employ, eqs. (12, 14) are true only for the complete correlation function. However we have used them for either the connected or the disconnected part separately. The best way to circumvent this problem is to make simulations at different strange quark masses; differentiation as in eq. (8) would then give directly $m_s \langle N | \bar{s}s | N \rangle$, the strange content of the nucleon. However this calculation is not feasible at the present time. This should be the ultimate goal of lattice simulations, as other recent (non-lattice) theoretical results, [6], have hinted that perhaps the strange quark content of the nucleon is not as large as supposed, previous results being explained by a combination of factors, such as $\Sigma \neq \sigma_{\pi N}$ and higher order corrections to first order perturbation theory. (Indeed there are already tantalising lattice indications that this may be so, [16, 17].)

In conclusion we would just like to say that lattice results at present are generally in qualitative agreement with other theoretical and experimental results. However much improvement in the calculations is required to be able to make quantitative predictions.

Acknowledgements

This work was supported in part by the Deutsche Forschungsgemeinschaft. The numerical computations were performed on the Cray Y-MP in Jülich with time granted by the Scientific Council of the HLRZ. We wish to thank both institutions for their support.

References

- [1] J. Ashman et. al., *Nucl. Phys.* **B328** (1989) 1.
- [2] For a review see e.g. G. Höhler, *Landolt-Börnstein* Vol.1/9b part 2, editor H. Schopper (Springer, Berlin 1983).
- [3] T.-P. Cheng and L.-F. Li, *Gauge Theory of Elementary Particles* Oxford University Press (1988, reprinted).
- [4] T.-P. Cheng and R. Dashen, *Phys. Rev. Lett.* **26** (1971) 594.
- [5] J. Gasser, H. Leutwyler, M. P. Locher and M. E. Sainio, *Phys. Lett.* **B213** (1988) 85.
- [6] J. Gasser, H. Leutwyler and M. E. Sainio, *Phys. Lett.* **B253** (1991) 252.
- [7] R. Gupta, C. F. Baillie, R. G. Brickner, G. W. Kilcup, A. Patel and S. R. Sharpe. *Phys. Rev.* **D44** (1991) 3272.
- [8] M. Creutz, *Quarks, Gluons and Lattices* Cambridge University Press (1988, reprinted).
- [9] R. Altmeyer, K. D. Born, M. Göckeler, R. Horsley, E. Laermann and G. Schierholz, *Nucl. Phys.* **B389** (1993) 445.
- [10] J. Smit, *Nucl. Phys.* **B(Proc. Supp.)B20** (1991) 542.
- [11] L. Maiani, G. Martinelli, M. L. Paciello and B. Taglienti, *Nucl. Phys.* **B293** (1987) 420.
- [12] K. Bitar, A. D. Kennedy, R. Horsley, S. Meyer and P. Rossi, *Nucl. Phys.* **B313** (1989) 348.
- [13] A. Patel, *Nucl. Phys.* **B(Proc. Supp.)B26** (1992) 350.
- [14] F. R. Brown, F. P. Butler, H. Chen, N. H. Christ, Z. Dhong, W. Schaffer, L. I. Unger and A. Vaccarino, *Phys. Rev. Lett.* **67** (1991) 1062.
- [15] C. Bernard, T. Blum, T. A. DeGrand, C. DeTar, S. Gottlieb, A. Krasnitz, R. L. Sugar and D. Toussaint, *Preprint*, IUHET-232, AZPH-TH/93-09, IPS Research Report No. 93-05, UCSBTH-93-09, UHHEP 93/2, 1993.
- [16] S. Güsen, K. Schilling, R. Sommer, K.-H. Mütter and A. Patel, *Phys. Lett.* **B212** (1988) 216.
- [17] R. Sommer, $\pi - N$ *Newsletter* Vol.4/5 editors Cutcosky, Höhler, Kluge, Nefkens (Karlsruhe, 1991).